arXiv:1203.4591v1 [math.FA] 20 Mar 2012

OSTROWSKI TYPE INEQUALITIES INVOLVING THE RIGHT CAPUTO FRACTIONAL DERIVATIVES BELONG TO L_p

MEHMET ZEKI SARIKAYA

Abstract. In this paper, we have established Ostrowski type inequalities involving the right Caputo fractional derivatives belong to L_p spaces $(1 \leq p \leq \infty)$ via the right Caputo fractional Taylor formula with integral remainder.

1. Introduction

In 1938, the classical integral inequality proved by A.M. Ostrowski as the following:

Theorem 1. *Let* $f : [a,b] \to \mathbb{R}$ *be continuous on* $[a,b]$ *and differentiable on* (a,b) *whose derivative* $f' : (a,b) \to \mathbb{R}$ *is bounded on* (a,b), *i.e.,* $\|f'\|_\infty = \underset{t\in(a,b)}{sup} |f'(t)| < +\infty$. *Then the following inequality holds:*

$$\left| f(x) - \frac{1}{b-a}\int_a^b f(u)du \right| \leq (b-a)\left[\frac{1}{4} + \frac{\left(x-\frac{a+b}{2}\right)^2}{(b-a)^2}\right]\|f'\|_\infty \tag{1.1}$$

for any $x \in [a,b]$. *The constant* $\frac{1}{4}$ *is the best possible.*

In [1], Anastassiou established a new Ostrowski inequality which holds higer order derivatives functions.

Theorem 2. *Let* $f \in C^{n+1}([a,b])$, $n \in \mathbb{N}$ *and* $x \in [a,b]$ *be fixed, such that* $f^{(k)}(0) = 0$, $k = 1, ..., n$. *Then it holds*

$$\left| f(x) - \frac{1}{b-a}\int_a^b f(u)du \right| \leq \frac{\left\|f^{(n+1)}\right\|_\infty}{(n+2)!}\left[\frac{(x-a)^{n+2} + (b-x)^{n+2}}{(b-a)}\right]. \tag{1.2}$$

Clearly inequality (1.2) generalizes inequality (1.1) for higher order derivatives of f.

We give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used throughout this paper.

Definition 1. *Let* $f \in L_1[a,b]$. *The Riemann-Liouville integrals* $J_{a+}^{\alpha}f$ *and* $J_{b-}^{\alpha}f$ *of order* $\alpha > 0$ *with* $a \geq 0$ *are defined by*

$$J_{a+}^{\alpha}f(x) = \frac{1}{\Gamma(\alpha)}\int_a^x (x-t)^{\alpha-1} f(t)dt, \quad x > a$$

and

$$J_{b-}^{\alpha}f(x) = \frac{1}{\Gamma(\alpha)}\int_x^b (t-x)^{\alpha-1} f(t)dt, \quad x < b$$

2000 *Mathematics Subject Classification.* 26D07, 26D10, 26D15, 47A63.

Key words and phrases. Ostrowski inequality, Riemann-Liouville fractional integral, right Caputo fractional derivatives.

respectively where $\Gamma(\alpha)=\int_0^\infty e^{-t}u^{\alpha-1}du$. *Here is* $J_{a+}^0 f(x)=J_{b-}^0 f(x)=f(x)$.

Definition 2. *Let* $f\in AC^m([a,b])$ *(*$f^{(m-1)}$ *is in* $AC([a,b])$*),* $m\in\mathbb{N}$, $m=[\alpha]$, $\alpha>0$ *(*$[.]$ *the ceiling of the number). The right Caputo fractional derivative of order* $\alpha>0$ *is defined by*

$$D_{b-}^{\alpha}f(x)=\frac{(-1)^m}{\Gamma(m-\alpha)}\int_x^b (t-x)^{m-\alpha-1}f^{(m)}(t)dt,\quad x\leq b.$$

If $\alpha=m\in\mathbb{N}$, *then*

$$D_{b-}^{m}f(x)=(-1)^m f^{(m)}(x),\quad \forall x\in[a,b].$$

If $x>b$, *we define* $D_{b-}^{\alpha}f(x)=0$.

Properties concerning this operator can be found ([10]-[13]). For some recent results connected with fractional integral inequalities see ([1]-[13]).

In order to prove our main results, we need the following theorem proved by Anastassiou in [3].

Theorem 3. *Let* $f\in AC^m([a,b])$, $x\in[a,b]$, $m=[\alpha]$, $\alpha>0$. *Then*

$$f(x)=\sum_{k=0}^{m-1}\frac{f^{(k)}(b)}{k!}(x-b)^k+\frac{1}{\Gamma(\alpha)}\int_x^b (t-x)^{m-\alpha-1}D_{b-}^{m}f(t)dt$$

the right Caputo fractional Taylor formula with integral remainder.

In [5], Anastassiou established general univariate right Caputo fractional Ostrowski inequalities with respect to $\|.\|_p$, $1\leq p\leq\infty$.

Theorem 4. *Let* $f\in AC^m([a,b])$, $m=[\alpha]$ *and* $f^{(k)}(b)=0$, $k=1,...,m-1$. *Then*

$$\left|f(b)-\frac{1}{b-a}\int_a^b f(u)du\right| \tag{1.3}$$

$$\leq\begin{cases}\frac{\left\|D_{b-}^{m}f\right\|_{\infty,[a,b]}}{\Gamma(\alpha+2)}(b-a)^{\alpha}, & \text{if } D_{b-}^{\alpha}f\in L_\infty([a,b]),\ \alpha>0\\ \frac{\left\|D_{b-}^{m}f\right\|_{L_1([a,b])}}{\Gamma(\alpha+1)}(b-a)^{\alpha-1} & \text{if } D_{b-}^{\alpha}f\in L_1([a,b]),\ \alpha\geq1\\ \frac{\left\|D_{b-}^{m}f\right\|_{L_q([a,b])}}{\Gamma(\alpha)(p(\alpha-1)+1)^{\frac{1}{p}}(\alpha+\frac{1}{p})}(b-a)^{\alpha-1+\frac{1}{p}} & \begin{array}{c}\text{if } D_{b-}^{\alpha}f\in L_q([a,b]),\\ p,q>1,\ \frac{1}{p}+\frac{1}{q}=1,\ \alpha>1-\frac{1}{p}\end{array}\end{cases}$$

The aim of this paper is to establish Ostrowski type inequalities involving the right Caputo fractional derivatives belong to L_p spaces ($1\leq p\leq\infty$) via the right Caputo fractional Taylor formula with integral remainder.

2. Main Results

We start with the following theorem:

Theorem 5. *Let* $f,g\in AC^m([a,b])$, $m=[\alpha]$, $\alpha>0$. *Assume* $f^{(k)}(b)=g^{(k)}(b)=0$, $k=1,...,m-1$ *and* $D_{b-}^{\alpha}f, D_{b-}^{\alpha}g\in L_\infty([a,b])$. *Then*

$$\left|2\int_a^b f(x)g(x)dx-\int_a^b (f(x)g(b)+g(x)f(b))dx\right| \tag{2.1}$$

$$\leq \left\|D_{b-}^{\alpha}f\right\|_\infty J_{a+}^{\alpha+1}|g(b)|+\left\|D_{b-}^{\alpha}g\right\|_\infty J_{a+}^{\alpha+1}|f(b)|.$$

Proof. Let $x \in [a,b]$. Using Theorem 3 and from the hypothesis of Theorem 5, we have the following identities

$$f(x) - f(b) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} D_{b-}^{\alpha} f(t)dt \tag{2.2}$$

$$g(x) - g(b) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} D_{b-}^{\alpha} g(t)dt. \tag{2.3}$$

Multiplying both sides of (2.2) and (2.3) by $g(x)$ and $f(x)$ respectively and adding the resulting identities, we have

$$\begin{aligned} & 2f(x)g(x) - f(b)g(x) - f(x)g(b) \\ = \; & \frac{g(x)}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} D_{b-}^{\alpha} f(t)dt + \frac{f(x)}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} D_{b-}^{\alpha} g(t)dt. \end{aligned}$$

Integrating the resulting inequality with respest to x over $[a,b]$ and using the properties of modulus, we obtain

$$\begin{aligned} & \left| 2\int_a^b f(x)g(x)dx - \int_a^b (f(b)g(x) + f(x)g(b))\, dx \right| \\ & \qquad (2.4) \\ \leq \; & \int_a^b \frac{|g(x)|}{\Gamma(\alpha)} \left(\int_x^b (t-x)^{\alpha-1} \left| D_{b-}^{\alpha} f(t) \right| dt \right) dx + \int_a^b \frac{|f(x)|}{\Gamma(\alpha)} \left(\int_x^b (t-x)^{\alpha-1} \left| D_{b-}^{\alpha} g(t) \right| dt \right) dx. \end{aligned}$$

Hence it holds

$$\begin{aligned} & \left| 2\int_a^b f(x)g(x)dx - \int_a^b (f(b)g(x) + f(x)g(b))\, dx \right| \\ \leq \; & \frac{\left\| D_{b-}^{\alpha} f \right\|_{\infty}}{\Gamma(\alpha)} \int_a^b |g(x)| \left(\int_x^b (t-x)^{\alpha-1} dt \right) dx + \frac{\left\| D_{b-}^{\alpha} g \right\|_{\infty}}{\Gamma(\alpha)} \int_a^b |f(x)| \left(\int_x^b (t-x)^{\alpha-1} dt \right) dx \\ = \; & \frac{\left\| D_{b-}^{\alpha} f \right\|_{\infty}}{\Gamma(\alpha+1)} \int_a^b (b-x)^{\alpha} |g(x)|\, dx + \frac{\left\| D_{b-}^{\alpha} g \right\|_{\infty}}{\Gamma(\alpha+1)} \int_a^b (b-x)^{\alpha} |f(x)|\, dx \\ = \; & \left\| D_{b-}^{\alpha} f \right\|_{\infty} J_{a+}^{\alpha+1} |g(b)| + \left\| D_{b-}^{\alpha} g \right\|_{\infty} J_{a+}^{\alpha+1} |f(b)| \end{aligned}$$

which the proof is completed. □

Remark 1. *If we take $g(x) = 1$ in Theorem 5, the inequality (2.1) reduces the first inequality in (1.3).*

Theorem 6. *Let $f, g \in AC^m([a,b])$, $m = [\alpha]$, $\alpha \geq 1$. Assume $f^{(k)}(b) = g^{(k)}(b) = 0$, $k = 1, ..., m-1$ and $D_{b-}^{\alpha} f, D_{b-}^{\alpha} g \in L_1([a,b])$. Then*

$$\begin{aligned} & \left| 2\int_a^b f(x)g(x)dx - \int_a^b (f(x)g(b) + g(x)f(b))\, dx \right| \\ \leq \; & \left\| D_{b-}^{\alpha} f \right\|_{L_1([a,b])} J_{a+}^{\alpha} |g(b)| + \left\| D_{b-}^{\alpha} g \right\|_{L_1([a,b])} J_{a+}^{\alpha} |f(b)|. \end{aligned} \tag{2.5}$$

Proof. From the inequality (2.4) of Theorem 5, we have again

$$
\begin{aligned}
& \left|2\int_a^b f(x)g(x)dx-\int_a^b \left(f(b)g(x)+f(x)g(b)\right)dx\right| \\
\leq\ & \int_a^b \frac{|g(x)|}{\Gamma(\alpha)}\left(\int_x^b (t-x)^{\alpha-1}\left|D_{b-}^{\alpha}f(t)\right|dt\right)dx+\int_a^b \frac{|f(x)|}{\Gamma(\alpha)}\left(\int_x^b (t-x)^{\alpha-1}\left|D_{b-}^{\alpha}g(t)\right|dt\right)dx \\
\leq\ & \int_a^b \frac{|g(x)|}{\Gamma(\alpha)}(b-x)^{\alpha-1}\left(\int_x^b \left|D_{b-}^{\alpha}f(t)\right|dt\right)dx+\int_a^b \frac{|f(x)|}{\Gamma(\alpha)}(b-x)^{\alpha-1}\left(\int_x^b \left|D_{b-}^{\alpha}g(t)\right|dt\right)dx \\
=\ & \left\|D_{b-}^{\alpha}f\right\|_{L_1([a,b])}\int_a^b \frac{|g(x)|}{\Gamma(\alpha)}(b-x)^{\alpha-1}dx+\left\|D_{b-}^{\alpha}g\right\|_{L_1([a,b])}\int_a^b \frac{|f(x)|}{\Gamma(\alpha)}(b-x)^{\alpha-1}dx \\
=\ & \left\|D_{b-}^{\alpha}f\right\|_{L_1([a,b])}J_{a+}^{\alpha}|g(b)|+\left\|D_{b-}^{\alpha}g\right\|_{L_1([a,b])}J_{a+}^{\alpha}|f(b)|
\end{aligned}
$$

which completes the proof. □

Remark 2. *If we take $g(x)=1$ in Theorem 6, the inequality (2.5) reduces the second inequality in (1.3).*

Theorem 7. *Let $f,g\in AC^m([a,b])$, $m=[\alpha]$, $\alpha>0$, $p,q>1$, $\frac{1}{p}+\frac{1}{q}=1$, $\alpha>1-\frac{1}{p}$. Assume $f^{(k)}(b)=g^{(k)}(b)=0$, $k=1,...,m-1$ and $D_{b-}^{\alpha}f, D_{b-}^{\alpha}g\in L_q([a,b])$. Then*

$$
\begin{aligned}
& \left|2\int_a^b f(x)g(x)dx-\int_a^b \left(f(x)g(b)+g(x)f(b)\right)dx\right| \\
\leq\ & \Gamma\left(\alpha+\frac{1}{p}\right)\left(\left\|D_{b-}^{\alpha}f\right\|_{L_q([a,b])}J_{a+}^{\alpha+\frac{1}{p}}|g(b)|+\left\|D_{b-}^{\alpha}g\right\|_{L_q([a,b])}J_{a+}^{\alpha+\frac{1}{p}}|f(b)|\right).
\end{aligned}
\tag{2.6}
$$

Proof. From the inequality (2.4) of Theorem 5, we have again

$$
\begin{aligned}
& \left|2\int_a^b f(x)g(x)dx-\int_a^b \left(f(b)g(x)+f(x)g(b)\right)dx\right| \\
\leq\ & \int_a^b \frac{|g(x)|}{\Gamma(\alpha)}\left(\int_x^b (t-x)^{\alpha-1}\left|D_{b-}^{\alpha}f(t)\right|dt\right)dx+\int_a^b \frac{|f(x)|}{\Gamma(\alpha)}\left(\int_x^b (t-x)^{\alpha-1}\left|D_{b-}^{\alpha}g(t)\right|dt\right)dx.
\end{aligned}
$$

Using the Hölder's integral inequality, we obtain

$$\begin{aligned}
&\left|2\int_a^b f(x)g(x)dx-\int_a^b\left(f(b)g(x)+f(x)g(b)\right)dx\right|\\
\leq\ &\int_a^b\frac{|g(x)|}{\Gamma(\alpha)}\left(\int_x^b(t-x)^{(\alpha-1)p}dt\right)^{\frac{1}{p}}\left(\int_x^b\left|D_{b-}^{\alpha}f(t)\right|^q dt\right)^{\frac{1}{q}}dx\\
&+\int_a^b\frac{|f(x)|}{\Gamma(\alpha)}\left(\int_x^b(t-x)^{(\alpha-1)p}dt\right)^{\frac{1}{p}}\left(\int_x^b\left|D_{b-}^{\alpha}g(t)\right|^q dt\right)^{\frac{1}{q}}dx\\
=\ &\frac{\left\|D_{b-}^{\alpha}f\right\|_{L_q([a,b])}}{\Gamma(\alpha)}\int_a^b|g(x)|\left(\int_x^b(t-x)^{(\alpha-1)p}dt\right)^{\frac{1}{p}}dx\\
&+\frac{\left\|D_{b-}^{\alpha}g\right\|_{L_q([a,b])}}{\Gamma(\alpha)}\int_a^b|f(x)|\left(\int_x^b(t-x)^{(\alpha-1)p}dt\right)^{\frac{1}{p}}
\end{aligned}$$

By simple computation

$$\left(\int_x^b(t-x)^{(\alpha-1)p}dt\right)^{\frac{1}{p}}=\frac{(b-x)^{\alpha-1+\frac{1}{p}}}{(p(\alpha-1)+1)^{\frac{1}{p}}}.$$

This last equality is substituted the above, then we have the conclusion. □

Remark 3. *If we take $g(x)=1$ in Theorem 7, the inequality (2.6) reduces the third inequality in (1.3).*

References

[1] G.A. Anastassiou, *Ostrowski type inequalities*, Proc. AMS 123 (1995), 3775-3781.
[2] G.A. Anastassiou, *Fractional Differentiation Inequalities*, Research Monograph, Springer, New York, 2009.
[3] G.A. Anastassiou, *On Right Fractional Calculus*, Chaos, Solitons and Fractals, 42 (2009), 365-376.
[4] G.A. Anastassiou, *Univariate right fractional Ostrowski inequalities*, CUBO, accepted, 2011.
[5] G.A. Anastassiou, *Advances on Fractional Inequalities*, Springer, New York, 2011.
[6] S. Belarbi and Z. Dahmani, *On some new fractional integral inequalities*, J. Ineq. Pure and Appl. Math., 10(3) (2009), Art. 86.
[7] Z. Dahmani, *New inequalities in fractional integrals*, International Journal of Nonlinear Scinece, 9(4) (2010), 493-497.
[8] Z. Dahmani, L. Tabharit, S. Taf, *Some fractional integral inequalities*, Nonl. Sci. Lett. A, 1(2) (2010), 155-160.
[9] Z. Dahmani, L. Tabharit, S. Taf, *New generalizations of Gruss inequality usin Riemann-Liouville fractional integrals*, Bull. Math. Anal. Appl., 2(3) (2010), 93-99.
[10] R. Gorenflo, F. Mainardi, *Fractional calculus: integral and differential equations of fractional order*, Springer Verlag, Wien (1997), 223-276.
[11] S. Miller and B. Ross, *An introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons, USA, 1993, p.2.
[12] I. Podlubni, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
[13] S. G. Samko, A. A Kilbas, O. I. Marichev, *Fractional Integrals and Derivatives Theory and Application*, Gordan and Breach Science, New York, 1993.

E-mail address: `sarikayamz@gmail.com`